

**Natural Events Policy Documentation  
of a Natural Event Due to High Winds on 18-Mar-2004  
Kennewick, WA**

**Benton Clean Air Authority**

September 13, 2004

|   |    |
|---|----|
| OVERVIEW .....  | 2  |
| NATURAL EVENTS POLICY .....   | 2  |
| WASHINGTON STATE'S COLUMBIA PLATEAU .....                               | 2  |
| WINDBLOWN DUST NATURAL EVENTS ACTION PLAN .....                         | 2  |
| Definition of High Wind Event in NEAP .....                             | 3  |
| GENERAL DESCRIPTION OF AREA .....                                       | 4  |
| EVALUATION OF 18-MAR-2004 EVENT .....                                   | 6  |
| PM <sub>10</sub> Data.....  | 6  |
| High Wind Event Analysis .....  | 7  |
| Landscape Stability Conditions in the PM <sub>10</sub> Source Area..... | 15 |
| Agricultural BACM Assessment .....                                      | 17 |
| SUMMARY AND CONCLUSIONS .....   | 19 |
| ABBREVIATIONS AND ACRONYMS .....  | 20 |
| REFERENCES.....   | 21 |
| APPENDICIES .....   | 22 |

## **OVERVIEW**

On 18-Mar-2004 an exceedance of the primary 24-hour PM<sub>10</sub> National Ambient Air Quality Standard (NAAQS) was recorded at Kennewick, WA. PM<sub>10</sub> State/Local Air Monitoring Site (SLAMS). The concentration was officially reported as 301 µg/m<sup>3</sup> at standard atmospheric conditions. The primary contributor to the exceedance was identified as fallowed and planted wheat fields, vulnerable irrigated agricultural fields, and to a lesser extent, localized urban areas, which were subjected to high wind speeds. An exact quantification of the source contributions is not available. The Benton Clean Air Authority (BCAA) believes that the 18-Mar-2004 event is a “natural event” in accordance with the EPA Natural Events Policy.

## **NATURAL EVENTS POLICY**

The Natural Events Policy (NEP) was issued in May 1996 to provide an avenue of response to PM<sub>10</sub> air quality data that are due to uncontrollable natural events. Under NEP provisions, PM<sub>10</sub> attributable to a natural event can be excluded from an attainment or non-attainment decision. The NEP is applicable when PM<sub>10</sub> data is due to uncontrollable natural events and the dust originates from non-anthropogenic sources or from contributing anthropogenic sources controlled with best available control measures (BACM)

The two basic requirements of the NEP are:

- 1) The states must develop a Natural Events Action Plan (NEAP) to deal with future PM<sub>10</sub> NAAQS exceedances.
- 2) The states must also establish a clear and casual relationship between the observed natural event and the observed exceedance and document the event.

## **WASHINGTON STATE'S COLUMBIA PLATEAU WINDBLOWN DUST NATURAL EVENTS ACTION PLAN**

Washington State's Natural Events Action Plan (NEAP) to address PM<sub>10</sub> from natural events occurring in the Columbia Plateau region of eastern Washington was a result of a large number of PM<sub>10</sub> NAAQS exceedances in this region in the period from the late 1980's and early 1990's. Agricultural fields upwind of PM<sub>10</sub> monitoring sites were identified as the principal sources of windblown dust. The Washington State Department of Ecology's Air Program division developed the initial NEAP in 1998 and updated the document in 2003.

The NEAP has several purposes:

- Development of procedures for taking appropriate, reasonable measures to safeguard public health when natural events occur.
- Responsibility to assure that emission controls are applied to sources that contribute to exceedances of the PM<sub>10</sub> NAAQS, when those controls will result in fewer violations of the standards. Emission controls include BACM development and implementation.
- Authorization for documentation to be submitted to request designation of an exceedance of the NAAQS for PM<sub>10</sub> as being the result of a natural event.

### **Definition of High Wind Event in NEAP**

The 2003 NEAP refined the definition of high wind event for Washington State in accordance with the provisions of the NEP allowing the states to determine this definition. This provision recognizes the multiple variables that affect the wind erosion processes that result in windblown dust and the generation and transport of PM<sub>10</sub>, which geographically differs. Following is the definition of a “high wind event” from pages A1-A4 of the Washington State Columbia Plateau Windblown Dust Natural Events Action Plan (Ref 11):

*"A high wind event occurs when the wind entrains and suspends dust to the extent that concentrations of PM<sub>10</sub> are elevated. This occurs when the average hourly wind speed at 10m is 18 miles per hour or greater for two or more hours [18+2]; or in excess of 13 [13+2] miles per hour for two or more hours when conditions of higher susceptibility to wind erosion exist (see attachment A1). A high wind event that exceeds PM<sub>10</sub> standard is a natural event."*

This definition recognizes the concept that the wind speed threshold for wind erosive processes on soil to cause elevated PM<sub>10</sub> concentrations in the air is variable. This variability depends on multiple variables related to soil characteristics, wind gustiness, soil surface residue cover, moisture content, and others. Attachment A1 to the Appendix A and of the Columbia Plateau NEAP documents the research and explains the logic behind this two-stage “high wind event” definition. The high wind event definition also necessarily includes the concept that the intensity of the wind event is a combination of wind speed and significant duration (sustained wind).

### **Relationship of High Wind Event Definition to Documentation**

The amount of detail in the event documentation required by the NEAP varies with the category (18 mph for ≥ 2 hr; or 13 mph for ≥ 2 hr under higher wind erosion susceptibility) of high wind event definition. For the “18 + 2” category the documentation

burden is less because of the more clear-cut association of the observed PM<sub>10</sub> data and the wind speed profile. The wind speed profile contains wind speed, wind direction, and duration and essentially is a data plot of wind speed and direction against a period of time. Precipitation preceding the day of the event is also part the “18 + 2” data set. The higher wind speed event more easily meets the “clear and causal” criteria of the Natural Events Policy.


The “13 + 2” category within the high wind event definition is more complex with respect to establishing the link between the wind event and elevated PM<sub>10</sub>. The meteorological phenomena driving the process are spatially and temporally more complex and the number, geographic distribution and complexity of the meteorological measurements needed to describe the event are greater. These complexities may require more sophisticated methodology to reveal the dynamics of the event. Regional scale meteorological modeling coupled with PM<sub>10</sub> generation (emission) and transport modeling may be necessary and other data may be needed to link an event to remote source areas.

## **GENERAL DESCRIPTION OF AREA**

Kennewick, Richland (including West Richland), and Pasco, known collectively as the Tri-Cities, are located in southeast Washington where the Yakima, Columbia, and Snake Rivers meet (Figure 1). The eastern half of the State of Washington lies in the rain shadow of the Cascade Mountains making the region a semi-arid desert. Average annual precipitation in the Tri-Cities region is about 6-7 inches with high rainfall intensity being very uncommon. Irrigated agriculture produces a wide diversity of crops including fruits, vegetables, alfalfa and potatoes. Dryland (non-irrigated) wheat production compliments the irrigated cropping systems. Large areas of non-agricultural range and desert lands complete the major land use areas of the region.

The Tri-Cities are located in an open-ended river basin partially bounded by low hills to the south and southwest. The terrain coupled with prevailing south and west winds limit local stagnant air pollution by ventilating the area. This coupling can also produce some extraordinary wind speeds and patterns. These winds can produce significant wind erosion events that can blanket the Tri-Cities region with dust from vulnerable agricultural fields and other areas. On rare occasions, usually during the fall and winter, strong winds can occur from the north and northwest sectors.

Figure 1: The Tri-Cities Area



## EVALUATION OF 18-MAR-2004 EVENT

This section describes the major factors that affected the occurrence of the windblown dust event and an exceedance of the PM<sub>10</sub> NAAQS on 18-Mar-2004 in Kennewick, WA. Analysis of the high wind event summarizes the circumstances and characteristics of the event. Best available control measures (BACM) are reviewed to demonstrate compliance with the BACM requirement of the Natural Events Policy. Landscape stability conditions are described to show what factors on the land and the activities taking place contributed to the observed windblown dust PM<sub>10</sub> NAAQS exceedance.

### PM<sub>10</sub> Data

The Kennewick PM<sub>10</sub> federal reference method (FRM) monitor operates on a 1- in - 1 day schedule. Tables A1 and A2 in Appendix A shows Kennewick PM<sub>10</sub> data for 2003 and data for January through 17-Mar-2004, respectively. The average PM<sub>10</sub> concentration for 2003 was 29.2 ug/m<sup>3</sup>. The recorded exceedances of the PM<sub>10</sub> NAAQS in 2003 were 186 ug/m<sup>3</sup> on 5-Mar-2003, 1438 ug/m<sup>3</sup> on 28-Oct-2003 and 164 ug/m<sup>3</sup> on 10-Nov-2003. All of the exceedances from 2003 have been documented as Natural Events. The average PM<sub>10</sub> concentration in the months prior to the 18-Mar-2004 exceedance was 20.3 ug/m<sup>3</sup> in January 2004 and 15.6 ug/m<sup>3</sup> in February 2003. These daily PM<sub>10</sub> values in Tables A1 and A2 (Appendix A) show that the days with 24-Hour PM<sub>10</sub> concentrations that exceed the NAAQS are relatively rare and are much higher than the majority of daily values and other maximums for the period. The annual average PM<sub>10</sub> concentration has not exceeded the annual NAAQS standard of 50 ug/m<sup>3</sup> in 18 years of monitoring at the Kennewick site.

Table 1 shows the occurrence of windblown PM<sub>10</sub> exceedances, which have been documented as natural events since the inception of the NEP in May 1996. The BCAA takes principal responsibility for high wind events and natural events documentation for exceedances that affect primarily Benton County. Documentation of larger regional events that affect a greater area of the Columbia plateau including Benton County is the principal responsibility of the Washington State Department of Ecology's Air Program.

**Table 1: History of Documented Windblown Dust Natural Events in Benton County (BCAA jurisdiction)**

| <b>YEAR</b> | <b>DATE</b>  | <b>CONCENTRATION<br/><math>\mu\text{g}/\text{m}^3</math></b> | <b>ACTION TAKEN</b>           |
|-------------|--------------|--|-------------------------------|
| 1999        | September 23 | 180  | Ecology NEP <sup>1</sup>      |
| 1999        | September 25 | 305  | Ecology NEP <sup>1</sup>      |
| 2000        | July 31      | 218  | BCAA NEP <sup>2</sup>         |
| 2001        | March 13     | 351  | BCAA NEP <sup>2</sup>         |
| 2001        | September 25 | 284  | Ecology NEP <sup>1</sup>      |
| 2001        | October 23   | 267  | BCAA NEP <sup>2</sup>         |
| 2002        | August 16    | 186  | BCAA NEP Pending <sup>2</sup> |
| 2003        | March 5      | 186  | BCAA NEP <sup>2</sup>         |
| 2003        | October 28   | 1438   | Ecology NEP <sup>1</sup>      |
| 2003        | Nov 10       | 164  | BCAA NEP Pending <sup>2</sup> |

<sup>1</sup> Regional event with generalized dust storm conditions from a high wind event occurring in the intermountain region east of the Cascade Mountain range, which are documented by the Washington State Department of Ecology's Air Program.

<sup>2</sup> Dust storm conditions from a high wind event that affected primarily Benton County documented by the Benton Clean Air Authority.

## High Wind Event Analysis

### Synoptic Weather Pattern

The synoptic weather pattern provides the broad view of the weather systems that set up and drive the observed wind event. The positioning of high and low pressure areas with associated air mass circulation patterns and pressure gradients help in understanding the wind speeds, direction, duration, and shifting of winds that may occur during a wind event.

On 18-Mar-2004 a very organized, fast moving low-pressure system and corresponding cold front was beginning to make landfall on the Washington coast and a strong pressure gradient was developing across Washington State (Figure 2b). Because winds move from areas of higher pressure to lower pressure and the mass air movements are respectively counterclockwise and clockwise around lows and highs, strong winds began from the southwest towards the northeast at approximately 0900 (PST). During the course of the day, the low pressure system pushed closer to the weak highs, which were well established across the western United States. The resulting wind speeds were high enough to cause soil particles to become airborne and generate PM<sub>10</sub> emissions. These high, sustained winds continued until about 2200 (PST). By 19-Mar-2004 the low pressure system moved more inland over the north-central United States and high-pressure began to build in the Tri-Cities area (Figure 2c).

Figure 2 a-c: Synoptic Weather Maps

Figure 2a: 17-Mar-2004




Figure 2b: 18-Mar-2004





Figure 2c: 19-Mar-2004





### Wind Data in the PM<sub>10</sub> Source Area and the Urban Receptor Area

The following wind data shows in detail the manifestation of the larger scale synoptic circulation depicted in Figure 2. The wind direction for both urban and rural areas remained southwesterly for the duration of the highest wind speeds. Also for both areas, the wind speeds were more than sufficient for exceeding wind erosion thresholds and the wind travel (miles) was more than sufficient to accomplish transport of entrained PM<sub>10</sub> generated in the wind erosion process.

Wind data was analyzed from the Public Agricultural Weather Station (PAWS) meteorological (MET) network in locations in Benton County (Figure 3a). Figures 3b and 3c show wind data from rural and urban MET stations to show the general trends of wind speeds and directions in these two areas from 2000 (PST) 17-Mar-2004 through 0000 (PST) on 19-Mar-2004.

In the rural PM<sub>10</sub> source area of the Horse Heaven Hills dryland wheat growing area the wind speeds were above 18 mph for an extended period at Alderdale, Horrigan and Station 4 (Figure 3b) in a steady southwesterly direction. The [18+2] criteria was also met at the Benton City and WSU Tri-Cities urban PAWS stations (Figure 3c). The BCAA has observed this common or typical pattern on numerous dates for previous PM<sub>10</sub> NAAQS exceedances. The urban and rural PM<sub>10</sub> source area meets the wind data standard of [18+2] miles per hour for two or more hours.

Figure 3a: Benton County Area PAWS Stations




Figure 3b: Rural Wind Data (2000 PST 17-Mar-2004 through 0000 PST 19-Mar-2004)

|      | <b>Alderdale</b> |                | <b>Horrigan</b> |                | <b>Gramling</b> |                |
|------|------------------|----------------|-----------------|----------------|-----------------|----------------|
| Time | Wind Speed       | Wind Direction | Wind Speed      | Wind Direction | Wind Speed      | Wind Direction |
| 2000 | 4.56             | 227.200        | 2.37            | 149.200        | 9.12            | 229.700        |
| 2100 | 6.02             | 212.400        | 3.66            | 180.500        | 11.08           | 219.900        |
| 2200 | 6.14             | 212.900        | 3.31            | 200.100        | 12.52           | 246.100        |
| 2300 | 4.55             | 213.500        | 4.40            | 159.400        | 15.70           | 286.100        |
| 0000 | 5.09             | 214.400        | 4.98            | 173.000        | 16.55           | 286.500        |
| 0100 | 5.01             | 209.600        | 7.61            | 167.100        | 17.83           | 290.200        |
| 0200 | 5.45             | 231.400        | 4.95            | 180.300        | 15.77           | 269.700        |
| 0300 | 4.06             | 222.700        | 1.55            | 161.700        | 14.31           | 252.800        |
| 0400 | 6.59             | 242.700        | 2.44            | 3.405          | 11.29           | 255.100        |
| 0500 | 3.67             | 257.300        | 1.94            | 0.718          | 12.67           | 239.300        |
| 0600 | 4.69             | 251.100        | 4.66            | 220.400        | 9.10            | 215.200        |
| 0700 | 5.35             | 235.200        | 9.78            | 231.200        | 10.55           | 236.400        |
| 0800 | 6.43             | 238.500        | 10.02           | 219.500        | 11.15           | 255.300        |
| 0900 | 7.71             | 227.300        | 13.86           | 217.200        | 18.12           | 274.400        |
| 1000 | 7.75             | 221.200        | 18.63           | 227.200        | 21.14           | 274.200        |
| 1100 | 11.48            | 223.800        | 22.41           | 236.600        | 25.79           | 275.000        |
| 1200 | 20.00            | 275.600        | 23.67           | 248.000        | 25.59           | 274.200        |
| 1300 | 23.76            | 257.200        | 24.50           | 248.800        | 30.15           | 278.800        |
| 1400 | 20.51            | 249.400        | 23.58           | 251.800        | 30.22           | 281.200        |
| 1500 | 17.56            | 262.900        | 21.18           | 254.300        | 28.50           | 283.300        |
| 1600 | 18.03            | 274.000        | 21.32           | 250.200        | 26.98           | 287.300        |
| 1700 | 21.94            | 274.900        | 23.15           | 255.900        | 25.99           | 290.500        |
| 1800 | 20.42            | 279.400        | 20.00           | 274.500        | 23.80           | 290.100        |
| 1900 | 18.19            | 274.600        | 18.12           | 277.100        | 18.07           | 291.600        |
| 2000 | 13.99            | 256.700        | 20.74           | 278.600        | 14.02           | 283.900        |
| 2100 | 11.06            | 260.500        | 20.54           | 277.700        | 23.42           | 289.100        |
| 2200 | 6.69             | 258.400        | 11.42           | 287.100        | 20.09           | 292.000        |
| 2300 | 4.97             | 236.300        | 5.46            | 272.300        | 13.01           | 285.200        |
| 0000 | 5.45             | 246.300        | 3.20            | 212.200        | 11.06           | 261.300        |

Figure 3c: Urban Wind Data (2000 PST 17-Mar-2004 through 0000 PST 19-Mar-2004)

| Time | Kennewick  |                | Badger Canyon |                | Benton City |                | WSU Tri-Cities |                |
|------|------------|----------------|---------------|----------------|-------------|----------------|----------------|----------------|
|      | Wind Speed | Wind Direction | Wind Speed    | Wind Direction | Wind Speed  | Wind Direction | Wind Speed     | Wind Direction |
| 2000 | 1.89       | 188.400        | 1.34          | 147.700        | .70         | 149.700        | 1.51           | 8.670          |
| 2100 | 3.88       | 208.200        | 8.33          | 167.700        | 1.38        | 278.200        | .98            | 6.771          |
| 2200 | 5.53       | 200.000        | 12.28         | 194.500        | 5.53        | 283.500        | 4.78           | 185.200        |
| 2300 | 8.61       | 217.500        | 11.50         | 204.600        | 5.31        | 290.000        | 6.71           | 210.700        |
| 0000 | 8.31       | 221.300        | 13.56         | 204.400        | 6.94        | 283.000        | 4.18           | 217.600        |
| 0100 | 7.54       | 221.100        | 13.85         | 205.000        | 7.34        | 266.600        | 4.89           | 208.400        |
| 0200 | 9.67       | 225.000        | 11.74         | 207.400        | 8.04        | 274.200        | 3.41           | 237.700        |
| 0300 | 6.74       | 211.900        | 11.55         | 199.600        | 5.78        | 288.500        | 4.29           | 204.400        |
| 0400 | 5.68       | 208.600        | 10.04         | 188.400        | 4.05        | 282.400        | 5.63           | 190.600        |
| 0500 | 3.90       | 198.800        | 10.90         | 178.300        | 1.20        | 43.490         | 3.98           | 188.200        |
| 0600 | 2.60       | 194.900        | 9.98          | 173.400        | 5.57        | 244.100        | .58            | 229.700        |
| 0700 | 2.39       | 194.300        | 4.81          | 281.800        | 5.30        | 254.000        | 1.06           | 45.810         |
| 0800 | 1.77       | 200.300        | 3.50          | 274.500        | 6.69        | 267.300        | 5.86           | 156.300        |
| 0900 | 2.21       | 58.620         | 7.32          | 199.500        | 9.07        | 294.600        | 5.11           | 133.400        |
| 1000 | 6.34       | 251.900        | 14.46         | 230.300        | 14.24       | 284.200        | 12.45          | 183.400        |
| 1100 | 10.26      | 256.200        | 17.67         | 234.600        | 15.10       | 298.500        | 16.26          | 210.300        |
| 1200 | 11.54      | 252.800        | 19.69         | 241.900        | 15.66       | 302.200        | 16.17          | 228.700        |
| 1300 | 12.53      | 243.500        | 21.52         | 237.500        | 18.84       | 301.300        | 19.08          | 217.600        |
| 1400 | 15.84      | 251.400        | 21.30         | 240.700        | 18.70       | 311.300        | 23.22          | 222.000        |
| 1500 | 15.93      | 261.500        | 21.34         | 255.700        | 16.46       | 312.200        | 21.81          | 228.100        |
| 1600 | 16.80      | 272.300        | 17.67         | 260.400        | 15.42       | 319.700        | 21.63          | 233.800        |
| 1700 | 13.24      | 281.300        | 12.54         | 268.500        | 12.49       | 329.400        | 17.25          | 241.700        |
| 1800 | 12.59      | 281.600        | 12.52         | 274.800        | 15.15       | 323.400        | 19.57          | 236.500        |
| 1900 | 11.46      | 268.400        | 13.22         | 243.100        | 20.22       | 325.900        | 19.39          | 235.500        |
| 2000 | 12.89      | 265.800        | 13.20         | 257.000        | 19.08       | 324.800        | 14.36          | 234.600        |
| 2100 | 14.05      | 272.000        | 14.13         | 257.000        | 12.59       | 331.900        | 13.00          | 222.600        |
| 2200 | 10.30      | 270.000        | 11.53         | 250.200        | 9.11        | 329.300        | 5.53           | 246.900        |
| 2300 | 6.90       | 254.600        | 9.30          | 235.700        | 8.67        | 316.500        | 6.46           | 227.900        |
| 0000 | 4.33       | 264.000        | 8.72          | 233.400        | 5.02        | 280.200        | 6.82           | 227.600        |

Because of the increased topographical surface roughness in the urban area, the hourly wind speeds were lower than the speeds in the rural areas. These wind speeds were sufficient to support transport of PM<sub>10</sub> across the urban area. In addition, the pattern of wind speeds and directions were consistent among four urban MET stations and were consistent with transport from the rural areas. Furthermore, because these wind speeds in the urban area were high enough to generate dust from construction sites with highly vulnerable soil surface conditions, urban fugitive dust was most likely a contributor to the exceedances.

As described, the necessary and sufficient conditions for wind conditions occurred on 18-Mar-2004 to cause an exceedance of the PM<sub>10</sub> NAAQS at the Kennewick monitoring station in the Tri-Cities. Since there are many irrigated and dryland fields that are located relatively close to the Kennewick monitoring site (within 5-40 miles), the wind speeds and duration necessary to transport dust from the fields to the population center are relatively low. This close proximity of agricultural fields to the populated areas (and the PM<sub>10</sub> SLAMS) makes the Tri-Cities distinct from other agricultural windblown dust situations, which typically have more remotely located source areas.

## Direct Observational Information on the High Wind Event

The arid climate in the wheat-producing land in the areas to the southwest of the Tri-Cities makes fallow farming a necessary water harvesting method. In the fallow system, one-half the land is allowed to collect a year's amount of precipitation without growing wheat plants on that land. The other half of the land has actively growing wheat, which is using the moisture collected in the previous year's fallow land plus the current year's precipitation.

The exposed soils in fallow areas were potentially susceptible to wind erosion on 18-Mar-2004. The degree of susceptibility depended on the dryland wheat yields in the previous year, which determines the amount of straw residue available for holding the soil against the wind. The overall contribution from just-planted areas would also vary according to available surface residue left on the surface after planting. The landscape stability of the rural PM<sub>10</sub> source area on 18-Mar-2004 was sufficiently low to allow wind erosion with the combined effects of below-normal antecedent precipitation and necessary agricultural operations disturbing the soil.

Recently planted spring wheat fields were vulnerable because crop cover was not sufficient to decrease wind speed at the soil surface below wind erosion threshold speeds and were subject to wind erosion and PM<sub>10</sub> emissions. Sustained winds blowing across these unprotected and unstabilized soil surfaces cause soil particles to become airborne and transported into the Tri-Cities urban area even when BACM and BMP are being used. Planted wheat fields where seedlings were emerging very sporadically across the field didn't prevent the soil from blowing. The saltation creep effect was clearly seen in several areas, as a number of the fields that were blowing were being impacted by the blowing dust and the movement of soil from adjacent fields. Fallow fields with sufficient residue from previous wheat harvest were also being subjected to high wind speeds. Planted wheat fields and fallowed fields with good residue on 18-Mar-2004 were holding very well and exhibited little or no soil movement.

Windblown dust can come from construction sites at wind speeds substantially less than that needed to generate and transport dust from the agricultural areas to the urban areas. Based primarily upon direct observational evidence by BCAA staff and generally confirmed by the BCAA complaint records, often wind speeds of 5-10 mph can produce dust from extremely disturbed and vulnerable soil surfaces present on construction sites. On construction sites in and around the urban areas, extreme soil disturbance, an almost total lack of vegetative residue, and frequent mechanical activity make these construction sites vulnerable to wind erosion. On 18-Mar-2004, construction site dust may have contributed somewhat, but the data and our observations show that dust from agricultural areas was the dominant contributor.

## **Landscape Stability Conditions in the PM<sub>10</sub> Source Area**

The largest source area for PM<sub>10</sub> in the 18-Mar-2004 event was the Horse Heaven Hills dryland wheat growing area and recently planted potato fields. Precipitation and its effects on wheat culture are the principal determinants of landscape stability. Precipitation on an event time-scale can modify the susceptibility of the soil surface to particle detachment. Such precipitation effects are operative in the period of a few days prior to the wind event that causes the wind erosion. Recent precipitation within a few days prior to the wind event can suppress the amount of PM<sub>10</sub> emissions and lack of precipitation could have the opposite effect. Sufficient precipitation can suppress emissions to the point that no exceedance occurs.

The months leading up to the exceedance date of 18-Mar-2004 had above average precipitation recorded at the Hanford Area which is approximately 25 miles northwest of the Tri-Cities area. Although above average total precipitation is shown in Table A3 in Appendix A for the months October 2002 – March 2004, daily precipitation measurements from three Public Agricultural Weather System (PAWS) stations in rural Benton County and four (PAWS) stations in the urban agricultural areas Benton County showed no precipitation for 14 consecutive days prior to the 18-Mar-2004 wind event. (Figure 4). Therefore, bare soil and low precipitation in the months previous to October 2002 created below average residue areas, which are highly vulnerable to wind erosion, received no stabilizing effects of precipitation for two weeks before the wind event.

**Figure 4: Precipitation Prior to 18-Mar-2004 Exceedance**

|             | Alderdale | Horrigan | Gramling | WSU<br>TC | CBC<br>Pasco | Badger<br>Canyon | Kennewick |
|-------------|-----------|----------|----------|-----------|--------------|------------------|-----------|
| 22-Feb-2004 | .00       | .00      | .00      | .00       | .00          | .00              | .00       |
| 23-Feb-2004 | .02       | .02      | .00      | .00       | .00          | .00              | .00       |
| 24-Feb-2004 | .00       | .00      | .03      | .00       | .00          | .00              | .00       |
| 25-Feb-2004 | .09       | .09      | .12      | .10       | .10          | .11              | .09       |
| 26-Feb-2004 | .03       | .04      | .07      | .04       | .03          | .07              | .03       |
| 27-Feb-2004 | .04       | .09      | .04      | .30       | .07          | .13              | .04       |
| 28-Feb-2004 | .00       | .00      | .03      | .23       | .00          | .04              | .02       |
| 29-Feb-2004 | .00       | .00      | .00      | .00       | .00          | .00              | .00       |
| 1-Mar-2004  | .00       | .00      | .00      | .00       | .00          | .00              | .00       |
| 2-Mar-2004  | .00       | .00      | .00      | .00       | .00          | .00              | .00       |
| 3-Mar-2004  | .10       | .20      | .19      | .06       | .05          | .07              | .00       |
| 4-Mar-2004  | .00       | .00      | .00      | .00       | .00          | .00              | .00       |
| 5-Mar-2004  | .00       | .00      | .00      | .00       | .00          | .00              | .00       |
| 6-Mar-2004  | .00       | .00      | .00      | .00       | .00          | .00              | .00       |
| 7-Mar-2004  | .00       | .00      | .00      | .00       | .00          | .00              | .00       |
| 8-Mar-2004  | .00       | .00      | .00      | .00       | .00          | .00              | .00       |
| 9-Mar-2004  | .00       | .00      | .00      | .00       | .00          | .00              | .00       |
| 10-Mar-2004 | .00       | .00      | .00      | .00       | .00          | .00              | .00       |
| 11-Mar-2004 | .00       | .00      | .00      | .00       | .00          | .00              | .00       |
| 12-Mar-2004 | .00       | .00      | .00      | .00       | .00          | .00              | .00       |
| 13-Mar-2004 | .00       | .00      | .00      | .00       | .00          | .00              | .00       |
| 14-Mar-2004 | .00       | .00      | .00      | .00       | .00          | .00              | .00       |
| 15-Mar-2004 | .00       | .00      | .00      | .00       | .00          | .00              | .00       |
| 16-Mar-2004 | .00       | .00      | .00      | .00       | .00          | .00              | .00       |
| 17-Mar-2004 | .00       | .00      | .00      | .00       | .00          | .00              | .00       |
| 18-Mar-2004 | .00       | .00      | .00      | .00       | .00          | .00              | .00       |



Most likely, the largest sources of dust on 18-Mar-2004 were from dryland fields disturbed by recent seeding of spring wheat fields and irrigated fields disturbed by recent seeding of potatoes. Spring wheat fields are particularly vulnerable to wind erosion during preparation for planting and following planting because of the disturbed soil condition. Any tillage that prepares the fields for planting and the planting operations themselves reduce surface residue. This reduction combined with an overall reduced supply of residue from low wheat yields in the previous crop cycle further increases susceptibility to wind erosion.

In summary, the combined effects of deficit and lack of rainfall for over 14 consecutive days (greater than 72 hours) prior to 18-Mar-2004 resulted in a vulnerable unstable landscape that was susceptible to wind erosion when the 18-Mar-2004 wind event occurred.

### **Agricultural BACM Assessment**

For agricultural sources, BACM is more commonly referred to as Best Management Practices (BMPs). A variety of management practices to control wind erosion and associated PM<sub>10</sub> emissions were one of the expected outcomes of the Columbia Plateau PM<sub>10</sub> Project. To qualify as a BMP, the practice must be proven to reduce wind erosion significantly below that which would occur with bare and tilled soil under similar weather conditions. Meteorological and climatological conditions strongly affect effective wind erosion or dust control on agricultural lands. Maintaining soil stability on agricultural fields is a problem in the Tri-Cities region principally during the most vulnerable times, such as crop planting and harvesting, or for other tillage operations that leave the soil vulnerable to wind erosion.

In the 2003 NEAP, Washington State found that BACM is implemented throughout the Columbia Plateau, which includes Benton County. Washington State evaluated BACM implementation for agricultural fields using Core 4 data. The Core 4 data shows 68% of total farmable acres of the Columbia Plateau are either part of a USDA conservation program, use one of several minimum till practices, or have 15 to 30% residue on the soil. Based on this evaluation, Washington State views these levels of wind erosion control as sufficient to fulfill BACM criterion of the Natural Events Policy.

In addition to currently implemented BACM there are on-going efforts to enhance wind erosion controls on the Horse Heaven Hills in Benton County. A \$65,000 grant from EPA has made it possible for the Washington Department of Ecology to contract with the Benton Conservation District to carry out a project as an extension of the Columbia Plateau Wind Erosion/Air Quality Project (formerly CP3). The goals of the project are:

- 1) To provide immediate, temporary treatment to critical areas.
- 2) To promote other options for longer term or permanent wind erosion control measures. Specifically, a new six-bale Newhouse straw mulcher was purchased to apply straw on highly erodible areas. During parts of the last two cropping cycles,

approximately 700 tons of surface residue was applied to protect against the occurrence of windblown dust from these areas. Conservation Reserve Program (CRP) acres have also increased over 100% since the latest USDA CRP signup, which brings the total CRP acres in the Horse Heaven Hills to 25,136 acres.

## **SUMMARY AND CONCLUSIONS**

From the evidence presented, the following conclusions can be drawn:

1. The Tri-Cities area and outlying agricultural areas were subjected to high wind speeds on 18-Mar-2004.
2. The urban and rural PM<sub>10</sub> source areas meet the wind data standard of [18+2] miles per hour for two or more hours.
3. Agricultural fields, which were highly susceptible to wind erosion during the 18-Mar- 2004 wind events, included fallowed wheat fields, recently planted spring wheat fields, and recently planted potato fields with insufficient crop residue were blowing most severely.
4. The combination of the wind event, which had the necessary wind speed, duration, and direction to generate and transport PM<sub>10</sub>, and the vulnerable landscape, caused the 18-Mar-2004 exceedance. Although the agricultural fields have BACM applied, there were certain conditions present including reduced residue due to extended drought and recent wheat planting – that allowed the wind to overcome BACM. In addition, any unprotected area of soil surface at construction sites or elsewhere in the landscape would have had potential to contribute to the exceedance.
5. In light of the previous statements and the acknowledgment that the Columbia Plateau is, in general, highly susceptible to high wind events, show that windblown dust is the most probable source of the PM<sub>10</sub>.
6. Based upon these conclusions, the BCAA considers the PM<sub>10</sub> concentration recorded on 18-Mar-2004 to have been caused by a high wind natural event and requests that the data for this date be flagged as such in the AIRS database.

## **ABBREVIATIONS AND ACRONYMS**

|                        |   |
|------------------------|---|
| BACM.....              | Best Available Control Measures   |
| BMP .....              | Best Management Practices   |
| SLAMS .....            | State and Local Air Monitoring Station  |
| EPA.....               | U.S. Environmental Protection Agency  |
| BCAA .....             | Benton Clean Air Authority  |
| BFWWCAPCA ..           | Benton Franklin Walla Walla Counties Air Pollution Control Authority,<br>renamed BCAA in 1995 |
| MET.....               | Meteorological  |
| NAAQS.....             | National Ambient Air Quality Standard   |
| PM <sub>10</sub> ..... | Particulate Matter, 10 microns in diameter  |
| Ecology .....          | Washington State Department of Ecology  |
| PST .....              | Pacific Standard Time   |
| NEP .....              | Natural Events Policy   |
| NEAP .....             | Natural Event Action Plan   |
| MOA .....              | Memorandum of Agreement   |
| RACM.....              | Reasonably Available Control Measures   |
| FDP .....              | BCAA Fugitive Dust Policy   |
| HMN .....              | Hanford Meteorological Network  |
| PAWS.....              | Public Agricultural Weather System operated by Washington State<br>University                 |
| CFR.....               | U.S. Code of Federal Regulations  |

## REFERENCES

- 1) Hoitink D. J. and K. W. Burk. 2003. Hanford Site Climatological Data Summary 2003 with Historical Data. Richland, WA: Pacific Northwest National Laboratory.
- 2) Pacific Northwest National Laboratory. 2004. Meteorological Database. Richland, WA: Pacific Northwest National Laboratory.
- 3) Lauer D. A. *et al.* 1998. Tri-Cities Area PM<sub>10</sub> Study Report. Richland, WA: Benton County Clean Air Authority.
- 4) United States Environmental Protection Agency. 1986. Guideline on the Identification and Use of Air Quality Data Affected by Exceptional Events. Office of Air Quality Planning and Standards.
- 5) Nichols, Mary D. 30-May-1996. Memorandum. "Areas affected by PM<sub>10</sub> Natural Events". United States Environmental Protection Agency.
- 6) National Oceanic and Atmospheric Administration. 2003. "Daily Weather Maps, Weekly Series March 15-21, 2004." Washington, DC: Climate Prediction Center.
- 7) Washington State University. 2004. Meteorological Database. Prosser, WA: Public Agricultural Weather System, Washington State University.
- 8) Papendick, Robert *et al.* 1998. Farming with the Wind: Best management Practices for Controlling Wind Erosion and Air Quality on Columbia Plateau Croplands. Pullman, WA: College of Agriculture and Home Economics, Washington State University.
- 9) Office of Air Quality Planning and Standards. 2003. Database. "Aerometric Information Retrieval System." Durham, NC: U. S. Environmental Protection Agency.
- 10) Pacific Northwest National Laboratories. 2003. Meteorological Database. Richland, WA: Pacific Northwest National Laboratory.
- 11) Washington State Department of Ecology. 2003 . Natural Events Action Plan. Lacey, WA: Washington State Department of Ecology Headquarters

## APPENDICES

### Appendix A

**Table A1: Daily PM<sub>10</sub> Concentration (ug/m<sup>3</sup>) for 2003**

| CA   | JAN  | FEB  | MAR   | APR  | MAY  | JUN  | JUL  | AUG  | SEP  | OCT    | NOV   | DEC  | MEAN  | MAX    | NO  |
|------|------|------|-------|------|------|------|------|------|------|--------|-------|------|-------|--------|-----|
| 1    | 8    | 5    | 22    | 10   | 11   | 16   | 23   | 32   | 36   | 48     | 40    | 8    | 21.6  | 48.0   | 12  |
| 2    | 12   | 5    | 17    | 6    | 21   | 25   | 18   | 26   | 58   | 49     | 26    | 18   | 23.4  | 58.0   | 12  |
| 3    | 7    |      |       | 32   | 28   | 24   | 24   | 21   |      | 56     | 17    | 23   | 25.8  | 56.0   | 9   |
| 4    | 5    | 13   | 11    | 11   | 33   | 25   | 21   | 47   | 75   | 53     | 22    | 10   | 27.2  | 75.0   | 12  |
| 5    | 6    | 21   | 186   | 10   | 14   |      | 20   | 45   | 49   | 47     | 40    | 15   | 41.2  | 186.0  | 11  |
| 6    | 13   | 26   | 126   | 4    | 13   | 34   |      | 19   | 86   | 59     |       | 10   | 39.0  | 126.0  | 10  |
| 7    | 11   | 27   | 21    | 10   | 17   | 27   |      | 18   |      | 19     | 39    | 12   | 20.1  | 39.0   | 10  |
| 8    | 12   | 20   | 7     | 16   | 18   | 36   | 36   | 16   | 15   | 84     | 37    | 13   | 25.8  | 84.0   | 12  |
| 9    | 11   | 22   | 33    | 13   | 20   | 27   | 27   | 80   | 13   | 30     | 32    | 9    | 26.4  | 80.0   | 12  |
| 10   |      | 22   | 17    | 17   | 17   | 33   |      | 15   | 29   | 14     | 164   | 5    | 33.3  | 164.0  | 10  |
| 11   | 14   | 20   | 34    | 7    | 15   | 19   | 42   | 14   | 5    | 16     | 41    | 12   | 19.9  | 42.0   | 12  |
| 12   | 14   | 18   |       | 11   | 13   | 28   | 69   | 16   | 16   | 138    | 38    | 19   | 34.5  | 138.0  | 11  |
| 13   | 14   | 16   | 60    | 9    | 16   | 37   | 25   | 30   | 22   |        | 40    | 11   | 25.5  | 60.0   | 11  |
| 14   | 13   | 22   | 19    |      | 34   | 15   | 22   | 34   | 27   | 14     | 50    | 3    | 23.0  | 50.0   | 11  |
| 15   | 9    | 17   |       | 9    | 39   | 17   | 30   | 62   |      | 13     |       | 8    | 22.7  | 62.0   | 9   |
| 16   | 12   | 4    |       |      | 8    | 28   | 25   | 41   | 54   | 1      | 18    | 10   | 20.1  | 54.0   | 10  |
| 17   |      | 4    |       | 11   | 9    | 34   | 22   | 17   | 14   | 15     | 31    | 12   | 16.9  | 34.0   | 10  |
| 18   | 10   |      | 12    | 7    | 9    | 65   | 29   |      | 24   | 23     |       | 13   | 21.3  | 65.0   | 9   |
| 19   | 12   | 15   | 10    | 15   | 14   | 50   | 33   | 50   | 81   |        | 10    | 15   | 27.7  | 81.0   | 11  |
| 20   | 6    | 14   | 36    | 14   | 18   | 17   | 23   | 41   | 20   | 22     | 7     | 18   | 19.7  | 41.0   | 12  |
| 21   |      | 9    | 6     | 13   | 16   | 20   | 25   | 54   |      | 14     | 16    | 11   | 18.4  | 54.0   | 10  |
| 22   | 5    | 5    | 9     | 17   | 27   | 9    | 43   | 40   | 33   | 42     | 14    |      | 22.2  | 43.0   | 11  |
| 23   | 12   | 10   | 11    | 16   | 72   | 16   | 36   | 21   | 36   | 22     | 13    | 11   | 23.0  | 72.0   | 12  |
| 24   | 19   | 14   | 11    | 15   | 24   | 19   | 39   | 16   | 43   | 22     | 2     | 10   | 19.5  | 43.0   | 12  |
| 25   | 21   | 27   |       | 4    | 13   | 22   | 34   |      | 40   | 24     | 37    | 2    | 22.4  | 40.0   | 10  |
| 26   | 4    | 25   | 19    | 2    | 9    | 30   | 35   | 31   | 45   | 27     | 12    | 7    | 20.5  | 45.0   | 12  |
| 27   | 6    | 25   | 16    | 3    | 10   | 22   | 29   | 17   | 41   |        | 12    |      | 18.1  | 41.0   | 10  |
| 28   | 9    | 19   | 13    |      | 22   | 22   | 37   | 24   | 38   | 1438   | 24    | 7    | 150.3 | 1438.0 | 11  |
| 29   | 10   |      | 14    | 5    | 21   | 32   | 40   | 55   | 61   | 40     | 7     | 10   | 26.8  | 61.0   | 11  |
| 30   | 11   |      | 20    | 5    | 28   | 43   | 44   | 45   | 55   | 126    |       |      | 41.9  | 126.0  | 9   |
| 31   | 6    |      | 29    |      | 17   |      | 35   | 40   |      | 39     |       | 22   | 26.9  | 40.0   | 7   |
| AVG  | 10.4 | 16.3 | 30.4  | 10.8 | 20.2 | 27.3 | 31.6 | 33.3 | 39.1 | 89.1   | 30.3  | 11.6 | 29.2  |        |     |
| MAX  | 21.0 | 27.0 | 186.0 | 32.0 | 72.0 | 65.0 | 69.0 | 80.0 | 86.0 | 1438.0 | 164.0 | 23.0 |       | 1438.0 |     |
| DAYS | 28.0 | 26.0 | 25.0  | 27.0 | 31.0 | 29.0 | 28.0 | 29.0 | 26.0 | 28.0   | 26.0  | 28.0 |       |        | 331 |

**Table A2: Daily PM<sub>10</sub> Concentration (ug/m<sup>3</sup>) for 2004**

| CA   | JAN  | FEB  | MAR   | APR | MAY | JUN | JUL | AUG | SEP | OCT | NOV | DEC | MEAN  | MAX   | NO |
|------|------|------|-------|-----|-----|-----|-----|-----|-----|-----|-----|-----|-------|-------|----|
| 1    | 16   | 7    |       |     |     |     |     |     |     |     |     |     | 11.5  | 16.0  | 2  |
| 2    | 10   |      | 13    |     |     |     |     |     |     |     |     |     | 11.5  | 13.0  | 2  |
| 3    | 16   | 9    | 11    |     |     |     |     |     |     |     |     |     | 12.0  | 16.0  | 3  |
| 4    | 24   | 20   | 6     |     |     |     |     |     |     |     |     |     | 16.7  | 24.0  | 3  |
| 5    | 40   | 27   | 30    |     |     |     |     |     |     |     |     |     | 32.3  | 40.0  | 3  |
| 6    | 14   | 17   | 10    |     |     |     |     |     |     |     |     |     | 13.7  | 17.0  | 3  |
| 7    | 17   | 15   | 14    |     |     |     |     |     |     |     |     |     | 15.3  | 17.0  | 3  |
| 8    | 30   | 17   | 20    |     |     |     |     |     |     |     |     |     | 22.3  | 30.0  | 3  |
| 9    | 50   | 20   | 22    |     |     |     |     |     |     |     |     |     | 30.7  | 50.0  | 3  |
| 10   | 44   | 23   | 30    |     |     |     |     |     |     |     |     |     | 32.3  | 44.0  | 3  |
| 11   | 40   | 20   | 23    |     |     |     |     |     |     |     |     |     | 27.7  | 40.0  | 3  |
| 12   | 15   | 16   | 19    |     |     |     |     |     |     |     |     |     | 16.7  | 19.0  | 3  |
| 13   | 16   | 25   | 14    |     |     |     |     |     |     |     |     |     | 18.3  | 25.0  | 3  |
| 14   | 23   | 29   | 7     |     |     |     |     |     |     |     |     |     | 19.7  | 29.0  | 3  |
| 15   | 23   | 22   |       |     |     |     |     |     |     |     |     |     | 22.5  | 23.0  | 2  |
| 16   | 10   | 12   | 19    |     |     |     |     |     |     |     |     |     | 13.7  | 19.0  | 3  |
| 17   | 17   | 12   | 19    |     |     |     |     |     |     |     |     |     | 16.0  | 19.0  | 3  |
| 18   | 19   | 13   | 301   |     |     |     |     |     |     |     |     |     | 111.0 | 301.0 | 3  |
| 19   | 13   | 13   |       |     |     |     |     |     |     |     |     |     | 13.0  | 13.0  | 2  |
| 20   | 20   | 15   |       |     |     |     |     |     |     |     |     |     | 17.5  | 20.0  | 2  |
| 21   | 21   | 19   |       |     |     |     |     |     |     |     |     |     | 20.0  | 21.0  | 2  |
| 22   | 31   | 14   |       |     |     |     |     |     |     |     |     |     | 22.5  | 31.0  | 2  |
| 23   | 20   |      |       |     |     |     |     |     |     |     |     |     | 20.0  | 20.0  | 1  |
| 24   | 9    | 15   |       |     |     |     |     |     |     |     |     |     | 12.0  | 15.0  | 2  |
| 25   | 4    | 9    |       |     |     |     |     |     |     |     |     |     | 6.5   | 9.0   | 2  |
| 26   | 6    | 9    |       |     |     |     |     |     |     |     |     |     | 7.5   | 9.0   | 2  |
| 27   | 15   | 7    |       |     |     |     |     |     |     |     |     |     | 11.0  | 15.0  | 2  |
| 28   |      | 7    |       |     |     |     |     |     |     |     |     |     | 7.0   | 7.0   | 1  |
| 29   | 5    | 9    |       |     |     |     |     |     |     |     |     |     | 7.0   | 9.0   | 2  |
| 30   | 22   |      |       |     |     |     |     |     |     |     |     |     | 22.0  | 22.0  | 1  |
| 31   |      |      |       |     |     |     |     |     |     |     |     |     | --    | --    | 0  |
| AVG  | 20.3 | 15.6 | 34.9  | --  | --  | --  | --  | --  | --  | --  | --  | --  | 21.8  |       |    |
| MAX  | 50.0 | 29.0 | 301.0 | --  | --  | --  | --  | --  | --  | --  | --  | --  |       | 301.0 |    |
| DAYS | 29.0 | 27.0 | 16.0  | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 | 0.0 |       |       | 72 |

**Table A3: Hanford Precipitation Data (10-1-2002 through 3-10-2004)**

|                          | Actual (in.) | Normal (in.) | Difference (in.) |
|--------------------------|--------------|--------------|------------------|
| October-02               | 0.12         | 0.49         | -0.37            |
| November-02              | 0.38         | 0.98         | -0.60            |
| December-02              | 2.36         | 1.11         | 1.25             |
| January-03               | 1.87         | 0.87         | 1.00             |
| February-03              | 0.82         | 0.68         | 0.14             |
| March-03                 | 0.26         | 0.58         | -0.32            |
| April-03                 | 2.23         | 0.44         | 1.79             |
| May-03                   | 0.08         | 0.55         | -0.47            |
| June-03                  | 0.00         | 0.41         | -0.41            |
| July-03                  | 0.00         | 0.27         | -0.27            |
| August-03                | 0.46         | 0.27         | 0.19             |
| September-03             | 0.24         | 0.31         | -0.09            |
| October-03               | 0.07         | 0.49         | -0.42            |
| November-03              | 0.15         | 0.98         | -0.83            |
| December-03              | 1.96         | 1.11         | 0.85             |
| January-04               | 2.12         | 0.87         | 1.25             |
| February-04              | 0.92         | 0.68         | 0.24             |
| March-04 <sup>1</sup>    | 0.15         | N/A          | N/A              |
| Totals (Oct '02-Mar '04) | 14.19        | 11.09        | 3.1              |

<sup>1</sup> Total precipitation before exceedance date of 3-18-2004